

Sum rules for total cross-sections of hadron photoproduction on ground state $1/2^+$ octet baryons

S. Dubnička,¹ A.Z. Dubničková,² and E. A. Kuraev³

¹*Inst. of Physics, Slovak Academy of Sciences, Dúbravská cesta 9, 845 11 Bratislava, Slovak Republic*

²*Dept. of Theor. Physics, Comenius University, 842 48 Bratislava, Slovak Republic*

³*Bogol'ubov Laboratory of Theoretical Physics, JINR, Dubna, Russia*

Sum rules are derived relating Dirac mean square radii and anomalous magnetic moments of various couples of the ground state $1/2^+$ octet baryons with the convergent integral of the difference of hadron photoproduction total cross-sections on the baryons. The chain of inequalities for corresponding total cross-sections has been found from those sum rules.

PACS numbers: 11.55.Hx, 13.60.Hb, 25.20.Lj

Keywords: sum rule, photoproduction, cross-section

Recently, the new sum rule has been derived [1]

$$\frac{1}{3}\langle r_{1p}^2 \rangle - \frac{\kappa_p^2}{4m_p^2} + \frac{\kappa_n^2}{4m_n^2} = \frac{2}{\pi^2\alpha} \int_{\omega_N}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma n \rightarrow X}(\omega)] \quad (1)$$

relating Dirac proton mean square radius $\langle r_{1p}^2 \rangle$ and anomalous magnetic moments of proton κ_p and neutron κ_n to the convergent integral over a difference of the total proton and neutron photoproduction cross-sections, in which a mutual cancellation of the rise of the corresponding cross-sections for $\omega \rightarrow \infty$ (ω is the photon energy in the laboratory frame), created by the Pomeron exchanges, was achieved. Using similar ideas the new Cabibbo-Radicati [2] like sum rules for various suitable couples of the members of the pseudoscalar meson nonet have been found in Ref. [3]. In this work, to be fascinated just by the very precise experimental satisfaction of the sum rule for a difference of proton and neutron total photoproduction cross-sections evaluating both sides of (1) (for more detail see [1]) and getting $(1.93 \pm 0.18)\text{mb}$ and $(1.92 \pm 0.32)\text{mb}$, respectively, we extend the method for a derivation of all possible sum rules for various suitable couples of the members of the ground state $1/2^+$ octet baryons.

Really, in the first place starting from a very high energy peripheral electroproduction process on baryon B

$$e^-(p_1) + B(p) \rightarrow e^-(p'_1) + X, \quad (2)$$

with the produced pure hadronic state X moving closely to the direction of the initial baryon, one comes (see [1]) to the Weizsäcker - Williams like relations, relating the difference of $\mathbf{q}^2$ -dependent (the photon transferred four-momentum $q^2 = -\mathbf{q}^2$) differential DIS cross-sections with integral over the difference of the total hadron photoproduction cross-sections on octet baryons.

Then utilizing analytic properties of parts of forward virtual Compton scattering amplitudes on octet baryons together with previous result one comes to the universal (compare with (1)) octet baryon sum rule

$$\begin{aligned} & \frac{1}{3} [F_{1B}(0)\langle r_{1B}^2 \rangle - F_{1B'}(0)\langle r_{1B'}^2 \rangle] - \left[\frac{\kappa_B^2}{4m_B^2} - \frac{\kappa_{B'}^2}{4m_{B'}^2} \right] = \\ & = \frac{2}{\pi^2\alpha} \int_{\omega_B}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma B \rightarrow X}(\omega) - \sigma_{tot}^{\gamma B' \rightarrow X}(\omega)] \end{aligned} \quad (3)$$

relating Dirac baryon mean square radii $\langle r_{1B}^2 \rangle, \langle r_{1B'}^2 \rangle$ and baryon anomalous magnetic moments $\kappa_B, \kappa_{B'}$ to the convergent integral over a difference of the total baryon photoproduction cross-sections. As there are 8 different members, $p, n, \Lambda, \Sigma^+, \Sigma^0, \Sigma^-, \Xi^0, \Xi^-$, of the ground state $1/2^+$ baryon octet, one can write down 28 different sum rules of the type (3). In order to evaluate the left-hand sides of them one uses the baryon masses and magnetic moments from Review of Particle Physics [4] (the Σ^0 magnetic moment is determined from the well known relation $\mu_{\Sigma^+} + \mu_{\Sigma^-} = 2\mu_{\Sigma^0}$) and Dirac baryon mean square radii $\langle r_{1B}^2 \rangle$ are calculated by means of the relation

$$\langle r_{1B}^2 \rangle = \langle r_{EB}^2 \rangle - \frac{3\kappa_B}{2m_B^2}, \quad (4)$$

where the baryon electric mean square radii $\langle r_{EB}^2 \rangle$ are calculated by Kubis and Meissner [5] to fourth order in relativistic baryon chiral perturbation theory, giving predictions for the Σ^- charge radius and $\Lambda - \Sigma^0$ transition moment in excellent agreement with the available experimental information.

As a result one gets numerically

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma n \rightarrow X}(\omega)] = 2.0415mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma n \rightarrow X}(\omega) \quad (5)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^+}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega)] = 2.0825mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) \quad (6)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^+}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega)] = 4.2654mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega) \quad (7)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega)] = 2.1829mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega) \quad (8)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Xi^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega)] = 1.5921mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega) \quad (9)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega)] = 1.6673mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) \quad (10)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega)] = -0.4158mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) \quad (11)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega)] = 1.6667mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) \quad (12)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega)] = 3.8496mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega) \quad (13)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega)] = 1.7259mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) \quad (14)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_p}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega)] = 3.3180mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega) \quad (15)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega)] = -0.3260mb, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) \quad (16)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega)] = -2.4573\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) \quad (17)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega)] = -0.3747\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) \quad (18)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega)] = 1.8082\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega) \quad (19)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega)] = -0.3156\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) \quad (20)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_n}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma n \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega)] = 1.2766\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma n \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega) \quad (21)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Lambda^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega)] = -2.0831\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) < \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) \quad (22)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Lambda^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega)] = -0.0006\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) \approx \sigma_{tot}^{\gamma \Sigma^0 \rightarrow X}(\omega) \quad (23)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Lambda^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega)] = 2.1823\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Sigma^- \rightarrow X}(\omega) \quad (24)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Lambda^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega)] = 0.0586\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) \quad (25)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Lambda^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega)] = 2.1823\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Lambda^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega) \quad (26)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^+}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega)] = 2.1417\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^0 \rightarrow X}(\omega) \quad (27)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^+}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) - \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega)] = 3.7338\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma \Sigma^+ \rightarrow X}(\omega) > \sigma_{tot}^{\gamma \Xi^- \rightarrow X}(\omega) \quad (28)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma\Sigma^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma\Xi^0 \rightarrow X}(\omega)] = 0.1168\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma\Sigma^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Xi^0 \rightarrow X}(\omega) \quad (29)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^0}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma\Sigma^0 \rightarrow X}(\omega) - \sigma_{tot}^{\gamma\Xi^- \rightarrow X}(\omega)] = 1.5732\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma\Sigma^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Xi^- \rightarrow X}(\omega) \quad (30)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^-}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma\Sigma^- \rightarrow X}(\omega) - \sigma_{tot}^{\gamma\Xi^0 \rightarrow X}(\omega)] = -2.1238\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma\Sigma^- \rightarrow X}(\omega) < \sigma_{tot}^{\gamma\Xi^0 \rightarrow X}(\omega) \quad (31)$$

$$\frac{2}{\pi^2\alpha} \int_{\omega_{\Sigma^-}}^{\infty} \frac{d\omega}{\omega} [\sigma_{tot}^{\gamma\Sigma^- \rightarrow X}(\omega) - \sigma_{tot}^{\gamma\Xi^- \rightarrow X}(\omega)] = -0.5316\text{mb}, \quad \text{then in averaged } \sigma_{tot}^{\gamma\Sigma^- \rightarrow X}(\omega) < \sigma_{tot}^{\gamma\Xi^- \rightarrow X}(\omega), \quad (32)$$

from where the following chain of inequalities

$$\sigma_{tot}^{\gamma\Sigma^+ \rightarrow X}(\omega) > \sigma_{tot}^{\gamma p \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Lambda^0 \rightarrow X}(\omega) \approx \sigma_{tot}^{\gamma\Sigma^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Xi^0 \rightarrow X}(\omega) > \sigma_{tot}^{\gamma n \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Xi^- \rightarrow X}(\omega) > \sigma_{tot}^{\gamma\Sigma^- \rightarrow X}(\omega) \quad (33)$$

for total cross-sections of hadron photoproduction on ground state $1/2^+$ octet baryons is found.

Experimental tests of the derived sum rules, as well as inequalities (33), could be practically carried out provided there exist data on total hadron photoproduction cross-sections on hyperons as a function of energy which, however, are missing up to present time. Nevertheless, the idea of intensive photon beams generated by the electron beams of linear e^+e^- colliders by using the process of the backward Compton scattering of laser light off the high energy electrons [6] is encouraging and one ex-

pects that a measurements of the total hadron photoproduction cross-sections on hyperons could be practically achievable in future.

The work was partly supported by Slovak Grant Agency for Sciences VEGA, Grant No. 2/4099/26 (S.D. and A.Z.D). A.Z. Dubničková would like to thank University di Trieste for warm hospitality at the early stage of this work and Professor N. Paver for numerous discussions.

-
- [1] E. Bartos, S.Dubnička and E. A. Kuraev, Phys. Rev. D70 (2004) 117901-1.
 - [2] N. Cabibbo, L. A. Radicati, Phys. Lett. 19 (1966) 697.
 - [3] S.Dubnička, A. Z. Dubničková, E.A. Kuraev, Phys.Rev. D74 (2006) 034023.

- [4] Review of Particle Physics, J. Phys. G 33 (2006).
- [5] B. Kubis and U.-G. Meissner, Eur. Phys. J. C 18 (2001) 747.
- [6] I.F. Ginzburg, G.L. Kotkin, V.G. Serbo and V.I. Telnov, JETP Lett. 34 (1981) 491.